

Estimate of dimension of Noether-Lefschetz locus for Beilinson-Hodge cycles on open complete intersections

M. Asakura and S. Saito

Contents

§0	Introduction
§1	Jacobian rings for open complete intersections
§2	Trivial part of cohomology of complements of hypersurfaces
§3	Hodge theoretic implication of generalized Jacobian rings
§4	Case of plane curves
§5	Beilinson's Tate conjecture
§6	Implication on injectivity of Chern class map for K_1 of surfaces
	References

0. Introduction

In his lectures in [G1], M. Green gives a lucid explanation how fruitful the infinitesimal method in Hodge theory is in various aspects of algebraic geometry. A significant idea is to use Koszul cohomology for Hodge-theoretic computations. The idea originates from Griffiths work [Gri] where the Poincaré residue representation of the cohomology of a hypersurface played a crucial role in proving the infinitesimal Torelli theorem for hypersurfaces. Since then many important applications of the idea have been made in different geometric problems such as the generic Torelli problem and the Noether-Lefschetz theorem for Hodge cycles and the study of algebraic cycles (see [G1, Lectures 7 and 8]).

In this paper we apply the method to study an analog of the Noether-Lefschetz theorem in the context of Beilinson's Hodge conjecture. Beilinson's Hodge conjecture and its Tate variant concern the regulator maps for open varieties (cf. [J1, Conjecture 8.5 and 8.6]). To be more precise we let U be a smooth variety over a field k of characteristic zero.

(0-1) (Hodge version) When $k = \mathbb{C}$ we have the regulator map from the higher Chow group to the singular cohomology of $U(\mathbb{C})$ (cf. [Bl] and [Sch])

$$\mathrm{reg}_U^q : CH^q(U, q) \otimes \mathbb{Q} \rightarrow H^q(U, \mathbb{Q}(q)) \cap F^q H^q(U, \mathbb{C})$$

where $\mathbb{Q}(q) = (2\pi\sqrt{-1})^q\mathbb{Q}$ and F^* denotes the Hodge filtration of the mixed Hodge structure on the singular cohomology defined by Deligne [D1]. Taking a smooth compactification $U \subset X$ with $Z = X \setminus U$, a simple normal crossing divisor on X , we have the following formula for the value of reg_U^q on decomposable elements in $CH^q(U, q)$;

$$reg_U^q(\{g_1, \dots, g_q\}) = \frac{dg_1}{g_1} \wedge \dots \wedge \frac{dg_q}{g_q} \in H^0(X, \Omega_X^q(\log Z)) = F^q H^q(U, \mathbb{C}),$$

where $\{g_1, \dots, g_q\} \in CH^q(U, q)$ is the products of $g_j \in CH^1(U, 1) = \Gamma(U, \mathcal{O}_{Zar}^*)$.

(0-2) (Tate version) When k is a finitely generated field over $\mathbb{Q}$ we have the étale regulator map from the higher Chow group to the ℓ -adic étale cohomology

$$reg_{et,U}^q : CH^q(U, q) \otimes \mathbb{Q}_\ell \rightarrow H_{et}^q(U_{\bar{k}}, \mathbb{Q}_\ell(j))^{Gal(\bar{k}/k)}$$

where $U_{\bar{k}} = U \times_k \bar{k}$ with $\bar{k}$, an algebraic closure of k .

Conjecture 0.1 *In case $k = \mathbb{C}$, reg_U^q is surjective.*

Conjecture 0.2 *In case k is finitely generated over $\mathbb{Q}$, $reg_{et,U}^q$ is surjective.*

We call them Beilinson's Hodge and Tate conjectures respectively. They are analogs of the Hodge and Tate conjectures for algebraic cycles on smooth projective varieties. The following are some remarks on the conjectures.

(i) The conjectures hold in case $q = 1$. They follow from the injectivity of Abel-Jacobi map (and its ℓ -adic variant) for divisors on smooth proper varieties.

(ii) If U is proper over k , we have

$$\begin{aligned} H^q(U, \mathbb{Q}(q)) \cap F^q H^q(U, \mathbb{C}) &= 0 \quad \text{in case } k = \mathbb{C}, \\ H_{et}^q(U_{\bar{k}}, \mathbb{Q}_\ell(q))^{Gal(\bar{k}/k)} &= 0 \quad \text{in case } k \text{ is finitely generated over } \mathbb{Q}, \end{aligned}$$

by the Hodge symmetry and by the reason of weight respectively. Thus Beilinson's conjectures are interesting only for non-proper varieties.

(iii) When X is a projective smooth surface and $U \subset X$ is the complement of a simple normal crossing divisor $Z \subset X$, then the surjectivity of reg_U^2 or $reg_{et,U}^2$ has an implication on the injectivity of the regulator maps for $CH^2(X, 1)$, which can be viewed as an analog of the Abel's theorem for K_1 of surfaces. The detail will be explained in §6.

(iv) As a naive generalization of Beilinson's Hodge conjecture, one may expect the surjectivity of the more general regulator map (in case $k = \mathbb{C}$)

$$reg_U^{p,q} : CH^q(U, 2q-p) \otimes \mathbb{Q} \rightarrow (2\pi\sqrt{-1})^q W_{2q} H^p(U, \mathbb{Q}) \cap F^q H^p(U, \mathbb{C})$$

where W_* denotes the weight filtration. Jannsen ([J1, 9.11]) has shown that the above map in case $p = 1$ and $q \geq 3$ is odd, is not surjective in general by using a theorem of Mumford [Mu], which implies the Abel-Jacobi for cycles of codimension ≥ 2 on smooth projective varieties is not injective even modulo torsion.

Before introducing the setup of the main result of this paper, we explain its background, the Noether-Lefschetz problem for Hodge cycles. Let $X \subset \mathbb{P}^n$ be a smooth projective variety over $\mathbb{C}$. Recall that the Hodge conjecture predicts that the space of Hodge cycles in codimension q on X :

$$F^0 H^{2q}(X, \mathbb{Q}(q)) := H^{2q}(X, \mathbb{Q}(q)) \cap F^q H^{2q}(X, \mathbb{C})$$

is generated by classes of algebraic subvarieties on X . One defines the space of trivial cycles in codimension q on X to be

$$H^{2q}(X, \mathbb{Q}(q))_{triv} := \text{Image}(H^{2q}(\mathbb{P}^n, \mathbb{Q}(q)) \rightarrow H^{2q}(X, \mathbb{Q}(q))) \subset F^0 H^{2q}(X, \mathbb{Q}(q)).$$

It is generated by the class of the section on X of a linear subspace of codimension q in $\mathbb{P}^n$. Now let S be a non-singular quasi-projective variety over $\mathbb{C}$ and assume that we are given $\mathcal{X} \hookrightarrow \mathbb{P}_S^n$, an algebraic family over S of smooth projective varieties. Let X_t be the fiber of $\mathcal{X}$ over $t \in S$. Then the Noether-Lefschetz locus for Hodge cycles in codimension q on $\mathcal{X}/S$ is defined to be

$$S_{NL}^q = \{t \in S \mid F^0 H^{2q}(X_t, \mathbb{Q}(q)) \neq H^{2q}(X_t, \mathbb{Q}(q))_{triv}\}.$$

It is the locus of such $t \in S$ that there are non-trivial Hodge cycles in codimension q on X_t and hence that the Hodge conjecture is non-trivial for X_t . One can prove S_{NL}^q is the union of countable number of (not necessarily proper) closed algebraic subsets of S (cf. [CDK]).

The celebrated theorem of Noether-Lefschetz affirms that in case $\mathcal{X}/S$ is the universal family of smooth surface of degree ≥ 4 in $\mathbb{P}^3$, $S_{NL}^1 \neq S$, which implies the Picard group of general surface of that kind is generated by the class of a hyperplane section.

In this paper we propose the following analog of the above problem in the context of Beilinson's Hodge conjecture. Assume that we are given schemes over $\mathbb{C}$

$$(0-3) \quad Y = \bigcup_{1 \leq j \leq s} Y_j \hookrightarrow \mathbb{P}^n \hookleftarrow X \hookleftarrow Z = \bigcup_{1 \leq j \leq s} Z_j, \quad V = \mathbb{P}^n \setminus Y, \quad U = X \setminus Z$$

where X is projective smooth, $Y_j \subset \mathbb{P}^n$ is a smooth hypersurface, Y is a simple normal crossing divisor on $\mathbb{P}^n$, $Z_j = X \cap Y_j$ intersecting transversally, and Z is a simple normal crossing divisor on X . We are interested in *the space of Beilinson-Hodge cycles in degree q on U* :

$$F^0 H^q(U, \mathbb{Q}(q)) := H^q(U, \mathbb{Q}(q)) \cap F^q H^q(U, \mathbb{C})$$

and the regulator map $reg_U^q : CH^q(U, q) \otimes \mathbb{Q} \rightarrow F^0 H^q(U, \mathbb{Q}(q))$. Since U is affine, $H^q(U, \mathbb{Q}(q)) = 0$ for $q > \dim(U)$ by the weak Lefschetz theorem. Thus we are interested only in case $1 \leq q \leq \dim(U)$. For a positive integer $q \neq n$ we put

$$H^q(U, \mathbb{Q}(q))_{triv} = \text{Image}(H^q(V, \mathbb{Q}(q)) \rightarrow H^q(U, \mathbb{Q}(q))).$$

We will define a subspace $CH^q(U, q)_{triv} \subset CH^q(U, q) \otimes \mathbb{Q}$ such that (cf. Definition 2.1)

$$H^q(U, \mathbb{Q}(q))_{triv} = reg_U^q(CH^q(U, q)_{triv})$$

and hence that $H^q(U, \mathbb{Q}(q))_{triv} \subset \text{Image}(reg_U^q) \subset F^0 H^q(U, \mathbb{Q}(q))$. We note that in case $\dim(X) > 1$ we have (cf. Lemma 2.3)

$$CH^q(U, q)_{triv} = CH^q(U, q)_{dec}$$

where $CH^q(U, q)_{dec} \subset CH^q(U, q) \otimes \mathbb{Q}$ is the subspace generated by decomposable elements, namely products of elements in $CH^1(U, 1) = \Gamma(U, \mathcal{O}_{S_{Zar}}^*)$. Now assume that we are given schemes over S , a non-singular quasi-projective variety over $\mathbb{C}$,

$$(0-4) \quad \mathcal{Y} = \bigcup_{1 \leq j \leq s} \mathcal{Y}_j \hookrightarrow \mathbb{P}_S^n \hookleftarrow \mathcal{X} \hookleftarrow \mathcal{Z} = \bigcup_{1 \leq j \leq s} \mathcal{Z}_j, \quad \mathcal{V} = \mathbb{P}_S^n \setminus \mathcal{Y}, \quad \mathcal{U} = \mathcal{X} \setminus \mathcal{Z}$$

whose fibers satisfies the same conditions as (0-3). Let U_t be the fiber of $\mathcal{U}$ over $t \in S$.

Definition 0.3 *For $1 \leq q \leq \dim(U_t)$ we define the Noether-Lefschetz locus for Beilinson-Hodge cycles in degree q on $\mathcal{U}/S$ to be*

$$S_{NL}^q = \{x \in S \mid F^0 H^q(U_t, \mathbb{Q}(q)) \neq H^q(U_t, \mathbb{Q}(q))_{triv}\}.$$

By definition S_{NL}^q is the locus of such $t \in S$ that there are non-trivial Beilinson-Hodge cycles in degree q on U_t and hence that Beilinson's Hodge conjecture is non-trivial for U_t . We note that a standard method in Hodge theory shows that S_{NL}^q is the union of countable number of (not necessarily proper) closed analytic subsets of S .

A main purpose of this paper is to study the Noether-Lefschetz locus in case the fibers of $\mathcal{X}/S$ are smooth complete intersection of multi-degree $(d_1, \dots, d_r)$ in $\mathbb{P}^n$. In this case the Lefschetz theory implies $S_{NL}^q = \emptyset$ unless $q = m := n - r = \dim(U_t)$ (cf. Lemma2.3). Thus we are interested only in $S_{NL} := S_{NL}^m$. In order state the main result we will introduce an invariant $c_S(\mathcal{X}, \mathcal{Z})$ in §3 that measures the “generality” of the family (0-4), or how many independent parameters S contains (cf. Definition3.2 and Lemma3.4). Assume that the fibers of $\mathcal{Y}_j \subset \mathbb{P}_S^n$ for $1 \leq j \leq s$ is a hypersurface of degree e_j and put

$$\delta_{min} = \min\{d_i, e_j \mid 1 \leq j \leq s, 1 \leq i \leq r\}.$$

Theorem 0.4 *For any irreducible component $E \subset S_{NL}$,*

$$\text{codim}_S(E) \geq \delta_{min}(n - r - 1) + \sum_{1 \leq i \leq r} d_i - c_S(\mathcal{X}, \mathcal{Z}) - n.$$

The original proof of the Noether-Lefschetz theorem was based on the study of monodromy action on cohomology of surfaces (cf. [D2]). The idea of improving the Noether-Lefschetz theorem by means of the infinitesimal method in the theory of variations of Hodge structures was introduced by Carlson, Green, Griffiths and Harris (cf. [CGGH]). By using the method closer analyses have been made on the Noether-Lefschetz locus for Hodge cycles on hypersurfaces, particularly on the problem to give a lower bound of codimensions of its irreducible components and to determine the components of maximal dimension (cf. [G4], [G5], [V] and [Ot]).

The proof of Theorem0.4 follows the same line of arguments as [G4] by using the variation of mixed Hodge structures arising from cohomology of the fibers of $\mathcal{U}/S$. A technical renovation is the results stated in §1 on generalized Jacobian rings, which give an algebraic description of the infinitesimal part of the variation of mixed Hodge structures. It is a natural generalization of the Poincaré residue representation of the cohomology of hypersurfaces in [Gri].

Having a result such as Theorem0.4, a natural question to ask is if the estimate in the theorem is optimal. In this paper we can give a positive answer to this question only in case the fibers of $\mathcal{X}/S$ are plane curves (cf. Theorem4.1). In case the fibers of $\mathcal{X}/S$ are of dimension > 1 , the estimate seems far from being optimal. Indeed in a forthcoming paper [AS2] we will show that the optimal estimate of codimensions of the Noether-Lefschetz locus for Beilinson's Hodge cycles on complements of three hyperplanes in surfaces of degree d in $\mathbb{P}^3$ is given explicitly by a quadratic polynomial in d .

Concerning Beilinson's Tate conjecture, we will show the following results. Let k be a finitely generated field over $\mathbb{Q}$. Let S be a quasi-projective variety over k and assume that we are given schemes over S which satisfy the same condition as (0-4). Assume that the fibers of $\mathcal{X}/S$ are smooth complete intersection of multi-degree $(d_1, \dots, d_r)$ in $\mathbb{P}^n$. Write $m = n - r$.

Theorem 0.5 *Assume $\sum_{1 \leq i \leq r} d_i \geq n + 1 + c_S(\mathcal{X}, \mathcal{Z})$.*

- (1) *Let $K = k(S)$ be the function field of S . For any a finite generated field L over K we have*

$$H_{et}^m(U_L, \mathbb{Q}_\ell(m))^{Gal(\bar{L}/L)} = reg_{et, U_L}^m(CH^m(U_L, m)_{triv} \otimes \mathbb{Q}_\ell),$$

where $reg_{et, U_L}^m : CH^m(U_L, m) \otimes \mathbb{Q}_\ell \rightarrow H_{et}^m(U_L, \mathbb{Q}_\ell(m))^{Gal(\bar{L}/L)}$ is the etale regulator map for $U_L = \mathcal{U} \times_S \text{Spec}(L)$ (cf. (0-2)).

- (2) *Assume that k is a finite extension of $\mathbb{Q}$ and $S(k) \neq \emptyset$. Let $\pi : S \rightarrow \mathbb{P}_k^N$ be a dominant quasi-finite morphism. There exist a subset $H \subset \mathbb{P}_k^N(k)$ such that:*

- (i) *For $\forall x \in S$ such that $\pi(x) \in H$ and for any subgroup $G \subset \text{Gal}(k(\bar{x})/k(x))$ of finite index we have*

$$H_{et}^m(U_{\bar{x}}, \mathbb{Q}_\ell(m))^G = reg_{et, U_x}^m(CH^m(U_x, m)_{triv} \otimes \mathbb{Q}_\ell),$$

where $reg_{et, U_x}^m : CH^m(U_x, m) \otimes \mathbb{Q}_\ell \rightarrow H_{et}^m(U_{\bar{x}}, \mathbb{Q}_\ell(m))^{Gal(k(\bar{x})/k(x))}$ is the etale regulator map for U_x , the fiber of $\mathcal{U}$ over $x \in S$, with $\bar{x}$, a geometric point over x .

- (ii) *Let Σ be any finite set of primes of k and let k_v be the completion of k at $v \in \Sigma$. Then the image of H in $\prod_{v \in \Sigma} \mathbb{P}_k^N(k_v)$ is dense.*

Now we explain how the paper is organized. In §1 we state the fundamental results on the generalized Jacobian rings, the duality theorem and the symmetrizer lemma. The proof is given in another paper [AS1]. It is based on the basic techniques to compute Koszul cohomology developed by M. Green ([G2] and [G3]). In §2 we define the trivial part of cohomology of complements of unions of hypersurface sections in smooth projective varieties, which is necessary to set up the Noether-Lefschetz problem for Beilinson-Hodge cycles. In §3 Hodge theoretic implications of the results in §1 are stated, which plays a crucial role in the proof of Theorem0.4 given in this section. The case of plane curves is treated in §4, where the estimate in Theorem0.4 is shown to be optimal. Theorem0.5 is proven in §5 by using the results in §2. In §6 we explain an implication of the Beilinson's conjectures on the injectivity of Chern class maps for K_1 of surfaces.

1. Jacobian rings for open complete intersection

The purpose of this section is to introduce Jacobian rings for open complete intersections and state their fundamental properties. Throughout the whole paper, we fix integers $r, s \geq 0$ with $r + s \geq 1$, $n \geq 2$ and $d_1, \dots, d_r, e_1, \dots, e_s \geq 1$. We put

$$\mathbf{d} = \sum_{i=1}^r d_i, \quad \mathbf{e} = \sum_{j=1}^s e_j, \quad \delta_{\min} = \min_{\substack{1 \leq i \leq r \\ 1 \leq j \leq s}} \{d_i, e_j\}, \quad d_{\max} = \max_{1 \leq i \leq r} \{d_i\}, \quad e_{\max} = \max_{1 \leq j \leq s} \{e_j\}.$$

We also fix a field k of characteristic zero. Let $P = k[X_0, \dots, X_n]$ be the polynomial ring over k in $n+1$ variables. Denote by $P^l \subset P$ the subspace of the homogeneous polynomials of degree l . Let A be a polynomial ring over P with indeterminants $\mu_1, \dots, \mu_r, \lambda_1, \dots, \lambda_s$. We use the multi-index notation

$$\mu^{\underline{a}} = \mu_1^{a_1} \cdots \mu_r^{a_r}, \quad \lambda^{\underline{b}} = \lambda_1^{b_1} \cdots \lambda_s^{b_s}, \quad \text{for } \underline{a} = (a_1, \dots, a_r) \in \mathbb{Z}_{\geq 0}^r, \quad \underline{b} = (b_1, \dots, b_s) \in \mathbb{Z}_{\geq 0}^s.$$

For $q \in \mathbb{Z}$ and $\ell \in \mathbb{Z}$, we write

$$A_q(\ell) = \bigoplus_{\mathbf{a}+\mathbf{b}=q} P^{\underline{ad}+\underline{be}+\ell} \cdot \mu^{\underline{a}} \lambda^{\underline{b}} \quad (\mathbf{a} = \sum_{i=1}^r a_i, \quad \mathbf{b} = \sum_{j=1}^s b_j, \quad \underline{ad} = \sum_{i=1}^r a_i d_i, \quad \underline{be} = \sum_{j=1}^s b_j e_j)$$

By convention $A_q(\ell) = 0$ if $q < 0$.

Definition 1.1 For $\underline{F} = (F_1, \dots, F_r)$, $\underline{G} = (G_1, \dots, G_s)$ with $F_i \in P^{d_i}$, $G_j \in P^{e_j}$, we define the **Jacobian ideal** $J(\underline{F}, \underline{G})$ to be the ideal of A generated by

$$\sum_{1 \leq i \leq r} \frac{\partial F_i}{\partial X_k} \mu_i + \sum_{1 \leq j \leq s} \frac{\partial G_j}{\partial X_k} \lambda_j, \quad F_i, \quad G_j \lambda_j \quad (1 \leq i \leq r, 1 \leq j \leq s, 0 \leq k \leq n).$$

The quotient ring $B = B(\underline{F}, \underline{G}) = A/J(\underline{F}, \underline{G})$ is called the **Jacobian ring of** $(\underline{F}, \underline{G})$. We denote

$$B_q(\ell) = B_q(\ell)(\underline{F}, \underline{G}) = A_q(\ell)/A_q(\ell) \cap J(\underline{F}, \underline{G}).$$

Definition 1.2 Suppose $n \geq r + 1$. Let $\mathbb{P}^n = \text{Proj } P$ be the projective space over k . Let $X \subset \mathbb{P}^n$ be defined by $F_1 = \dots = F_r = 0$ and let $Z_j \subset X$ be defined by $G_j = F_1 = \dots = F_r = 0$ for $1 \leq j \leq s$. We also call $B(\underline{F}, \underline{G})$ the Jacobian ring of the pair $(X, Z = \cup_{1 \leq j \leq s} Z_j)$ and denote $B(\underline{F}, \underline{G}) = B(X, Z)$ and $J(\underline{F}, \underline{G}) = J(X, Z)$.

In what follows we fix $\underline{F}$ and $\underline{G}$ as Definition 1.1 and assume the condition

(1-1) $F_i = 0$ ($1 \leq i \leq r$) and $G_j = 0$ ($1 \leq j \leq s$) intersect transversally in $\mathbb{P}^n$.

We mention three main theorems. The first main theorem concerns with the geometric meaning of Jacobian rings.

Theorem 1.3 Suppose $n \geq r + 1$. Let X and Z be as Definition 1.2.

(1) For integers $0 \leq q \leq n - r$ and $\ell \geq 0$ there is a natural isomorphism

$$\phi_{X,Z}^q : B_q(\mathbf{d} + \mathbf{e} - n - 1 + \ell) \xrightarrow{\cong} H^q(X, \Omega_X^{n-r-q}(\log Z)(\ell))_{\text{prim}}.$$

Here $\Omega_X^p(\log Z)$ is the sheaf of algebraic differential q -forms on X with logarithmic poles along Z and ‘prim’ means the primitive part:

$$H^q(X, \Omega_X^p(\log Z)(\ell))_{\text{prim}} = \begin{cases} \text{Coker}(H^q(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^q) \rightarrow H^q(X, \Omega_X^q)) & \text{if } q = p \text{ and } s = \ell = 0, \\ H^q(X, \Omega_X^p(\log Z)(\ell)) & \text{otherwise.} \end{cases}$$

(2) There is a natural map

$$\psi_{(X,Z)} : B_1(0) \longrightarrow H^1(X, T_X(-\log Z))_{\text{alg}} \subset H^1(X, T_X(-\log Z))$$

which is an isomorphism if $\dim(X) \geq 2$. Here $T_X(-\log Z)$ is the $\mathcal{O}_X$ -dual of $\Omega_X^1(\log Z)$ and the group on the right hand side is defined in Definition 1.4 below. The following map

$$H^1(X, T_X(-\log Z)) \otimes H^q(X, \Omega_X^p(\log Z)) \longrightarrow H^{q+1}(X, \Omega_X^{p-1}(\log Z)).$$

induced by the cup-product and the contraction $T_X(-\log Z) \otimes \Omega_X^p(\log Z) \rightarrow \Omega_X^{p-1}(\log Z)$ is compatible through $\psi_{(X,Z)}$ with the ring multiplication up to scalar.

Roughly speaking, the generalized Jacobian rings describe the infinitesimal part of the Hodge structures of open variety $X \setminus Z$, and the cup-product with Kodaira-Spencer class coincides with the ring multiplication up to non-zero scalar. This result was originally invented by P. Griffiths in case of hypersurfaces and generalized to complete intersections by Konno [K]. Our result is a further generalization.

Definition 1.4 Let the assumption be as in Theorem 1.3. We define $H^1(X, T_X(-\log Z))_{\text{alg}}$ to be the kernel of the composite map

$$H^1(X, T_X(-\log Z)) \rightarrow H^1(X, T_X) \rightarrow H^2(X, \mathcal{O}_X),$$

where the second map is induced by the cup product with the class $c_1(\mathcal{O}_X(1)) \in H^1(X, \Omega_X^1)$ and the contraction $T_X \otimes \Omega_X^1 \rightarrow \mathcal{O}_X$. It can be seen that

$$\dim_k(H^1(X, T_X(-\log Z))/H^1(X, T_X(-\log Z))_{\text{alg}}) = \begin{cases} 1 & \text{if } X \text{ is a K3 surface,} \\ 0 & \text{otherwise.} \end{cases}$$

The second main theorem is the duality theorem for the generalized Jacobian rings.

Theorem 1.5 Let the notation be as above.

(1) There is a natural map (called the trace map)

$$\tau : B_{n-r}(2(\mathbf{d} - n - 1) + \mathbf{e}) \rightarrow k.$$

Let

$$h_p(\ell) : B_p(\mathbf{d} - n - 1 + \ell) \rightarrow B_{n-r-p}(\mathbf{d} + \mathbf{e} - n - 1 - \ell)^*$$

be the map induced by the following pairing induced by the multiplication

$$B_p(\mathbf{d} - n - 1 + \ell) \otimes B_{n-r-p}(\mathbf{d} + \mathbf{e} - n - 1 - \ell) \rightarrow B_{n-r}(2(\mathbf{d} - n - 1) + \mathbf{e}) \xrightarrow{\tau} k.$$

When $r > n$ we define $h_p(\ell)$ to be the zero map by convention.

(2) The map $h_p(\ell)$ is an isomorphism in either of the following cases.

- (i) $s \geq 1$ and $p < n - r$ and $\ell < e_{\max}$.
- (ii) $s \geq 1$ and $0 \leq \ell \leq e_{\max}$ and $r + s \leq n$.
- (iii) $s = \ell = 0$ and either $n - r \geq 1$ or $n - r = p = 0$.

(3) The map $h_{n-r}(\ell)$ is injective if $s \geq 1$ and $\ell < e_{\max}$.

We have the following auxiliary result on the duality.

Theorem 1.6 Assume $n - r \geq 1$ and consider the composite map

$$\eta_{(X,Z)} : H^0(X, \Omega_X^{n-r}(\log Z)) \xrightarrow{(\phi_{X,Z}^0)^{-1}} B_0(\mathbf{d} + \mathbf{e} - n - 1) \xrightarrow{h_{n-r}(0)^*} B_{n-r}(\mathbf{d} - n - 1)^*$$

where the second map is the dual of $h_{n-r}(0)$. Then $\eta_{(X,Z)}$ is surjective and we have (cf. Definition 1.7 below)

$$\text{Ker}(\eta_{(X,Z)}) = \wedge_X^{n-r}(G_1, \dots, G_s).$$

Definition 1.7 Let $G_1, \dots, G_s$ be as in Definition 1.1 and let $Y_j \subset \mathbb{P}^n$ be the smooth hypersurface defined by $G_j = 0$. Let $X \subset \mathbb{P}^n$ be a smooth projective variety such that Y_j ($1 \leq j \leq s$) and X intersect transversally. Put $Z_j = X \cap Y_j$. Take an integer q with $0 \leq q \leq s - 1$. For integers $1 \leq j_1 < \dots < j_{q+1} \leq s$, let

$$\omega_X(j_1, \dots, j_{q+1}) \in H^0(X, \Omega_X^q(\log Z)) \quad (Z = \sum_{1 \leq j \leq s} Z_j)$$

be the restriction of

$$\sum_{\nu=1}^{q+1} (-1)^{\nu-1} e_{j_\nu} \frac{dG_{j_1}}{G_{j_1}} \wedge \dots \wedge \frac{d\widehat{G_{j_\nu}}}{G_{j_\nu}} \wedge \dots \wedge \frac{dG_{j_{q+1}}}{G_{j_{q+1}}} \in H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^q(\log Y))$$

where $Y = \sum_{1 \leq j \leq s} Y_j \subset \mathbb{P}^n$. We let

$$\wedge_X^q(G_1, \dots, G_s) \subset H^0(X, \Omega_X^q(\log Z))$$

be the subspace generated by $\omega_X(j_1, \dots, j_{q+1})$. For $1 \leq j_1 < \dots < j_q \leq s - 1$ we have

$$e_1^{q-1} \cdot \omega_X(1, j_1, \dots, j_q) = \frac{dg_{j_1}}{g_{j_1}} \wedge \dots \wedge \frac{dg_{j_q}}{g_{j_q}} \quad \text{with } g_j = (G_j^{e_1}/G_1^{e_j})|_X \in \Gamma(U, \mathcal{O}_U^*) \quad (U = X \setminus Z)$$

and $\omega_X(1, j_1, \dots, j_q)$ with $1 < j_1 < \dots < j_q \leq s$ form a basis of $\wedge_X^q(G_1, \dots, G_s)$.

Our last main theorem is the generalization of Donagi's symmetrizer lemma [Do] (see also [DG], [Na] and [N]) to the case of open complete intersections at higher degrees.

Theorem 1.8 Assume $s \geq 1$. Let $V \subset B_1(0)$ is a subspace of codimension $c \geq 0$. Then the Koszul complex

$$B_p(\ell) \otimes \wedge^{q+1} V \rightarrow B_{p+1}(\ell) \otimes \wedge^q V \rightarrow B_{p+2}(\ell) \otimes \wedge^{q-1} V$$

is exact if one of the following conditions is satisfied.

- (i) $p \geq 0$, $q = 0$ and $\delta_{\min} p + \ell \geq c$.
- (ii) $p \geq 0$, $q = 1$ and $\delta_{\min} p + \ell \geq 1 + c$ and $\delta_{\min}(p + 1) + \ell \geq d_{\max} + c$.
- (iii) $p \geq 0$, $\delta_{\min}(r + p) + \ell \geq \mathbf{d} + q + c$, $\mathbf{d} + e_{\max} - n - 1 > \ell \geq \mathbf{d} - n - 1$ and either $r + s \leq n + 2$ or $p \leq n - r - [q/2]$, where $[*]$ denotes the Gaussian symbol.

2. Trivial part of cohomology of complements of hypersurfaces

In this section we introduce some notions necessary to set up the Noether-Lefschetz problem for Beilinson-Hodge cycles on the complement of the union of hypersurface sections in a smooth projective variety. We fix the base field k which is either $\mathbb{C}$ or finitely generated over $\mathbb{Q}$. Assume that we are given schemes over k

$$(2-1) \quad Y = \bigcup_{1 \leq j \leq s} Y_j \hookrightarrow \mathbb{P}^n \hookleftarrow X \hookleftarrow Z = \bigcup_{1 \leq j \leq s} Z_j, \quad V = \mathbb{P}^n \setminus Y, \quad U = X \setminus Z$$

where X is projective smooth, $Y_j \subset \mathbb{P}^n$ is a smooth hypersurface of degree e_j , Y is a simple normal crossing divisor on $\mathbb{P}^n$, $Z_j = X \cap Y_j$ intersecting transversally, and Z is a simple normal crossing divisor on X . In what follows $CH^i(*, j)$ denotes the Bloch's higher Chow group (cf. [Bl]).

Definition 2.1 Assume $s \geq 2$.

- (1) Let $G_j \in H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(e_j))$ be a non-zero element defining $Y_j \subset \mathbb{P}^n$. For $1 \leq j \leq s-1$ put

$$g_j = (G_j^{e_s} / G_s^{e_j})|_U \in \Gamma(U, \mathcal{O}_U^*) = CH^1(U, 1).$$

Let $CH^1(U, 1)_{\text{triv}} \subset CH^1(U, 1) \otimes \mathbb{Q}$ be the subspace generated by k^* and g_j with $1 \leq j \leq s-1$.

- (2) For $q \geq 1$ let $CH^q(U, q)_{\text{dec}} \subset CH^q(U, q) \otimes \mathbb{Q}$ be the subspace generated by the products of elements in $CH^1(U, 1)$ and let

$$CH^q(U, q)_{\text{triv}} \subset CH^q(U, q)_{\text{dec}}$$

to be the subspace generated by the products of elements in $CH^1(U, 1)_{\text{triv}}$.

- (3) For $q \geq 1$ let $H^0(X, \Omega_X^q(\log Z))_{\text{triv}} \subset H^0(X, \Omega_X^q(\log Z))$ be the subspace generated by the wedge products of $dg_j/g_j \in H^0(X, \Omega_X^1(\log Z))$ with $1 \leq j \leq s-1$.

- (4) Let $H^p(U, q)$ denotes the singular cohomology $H^p(U, \mathbb{Q}(q))$ of $U(\mathbb{C})$ in case $k = \mathbb{C}$ (resp. the étale cohomology $H_{\text{ét}}^p(\bar{U}, \mathbb{Q}_\ell(q))$ in case k is finitely generated over $\mathbb{Q}$, where $\bar{U} = U \times_k \bar{k}$ with $\bar{k}$, an algebraic closure of k). For $q \geq 1$ let

$$H^q(U, q)_{\text{triv}} \subset H^q(U, q)$$

be the subspace generated by the image of $CH^q(U, q)_{\text{triv}}$ under the regulator map introduced in (0-1) and (0-2).

In case $s = 0, 1$ we put $M_{\text{triv}} = 0$ by convention for $M = CH^q(U, q)$, $H^0(X, \Omega_X^q(\log Z))$, $H^q(U, q)$.

Remark 2.2 In case $k = \mathbb{C}$ we have

$$H^q(U, \mathbb{Q}(q))_{triv} \otimes_{\mathbb{Q}} \mathbb{C} = H^0(X, \Omega_X^q(\log Z))_{triv}$$

under the identification $F^q H^q(U, \mathbb{C}) = H^0(X, \Omega_X^q(\log Z))$.

Lemma 2.3 Assume $s \geq 1$ and $q \geq 1$.

- (1) If $\dim(X) > 1$, $CH^q(U, q)_{dec} = CH^q(U, q)_{triv}$.
- (2) If $q \neq n$, $H^q(U, q)_{triv} = \text{Image}(H^q(V, q) \rightarrow H^q(U, q))$.
- (3) If $X \subset \mathbb{P}^n$ is a smooth complete intersection, $H^q(U, q) = H^q(U, q)_{triv}$ for $q \neq \dim(X)$.

Proof. To show Lemma2.3(1) it suffices to show $CH^1(U, 1)_{triv} = CH^1(U, 1) \otimes \mathbb{Q}$, which follows at once from the exact sequence

$$0 \rightarrow k^* \rightarrow CH^1(U, 1) \rightarrow \bigoplus_{1 \leq j \leq s} \mathbb{Z} \rightarrow CH^1(X)$$

where the second map is induced by the residue maps along Z_j and the last by the class of Z_j in $CH^1(X)$.

We show Lemma2.3(3). Since U is affine by the assumption $s \geq 1$, $H^q(U, q) = 0$ for $q > \dim(X)$ by the weak Lefschetz theorem. Thus we may assume that $1 \leq q < \dim(X)$ so that $\dim(X) > 1$. If $s = 1$, $Z = Z_1$ is a smooth complete intersection of dimension ≥ 1 and we have the exact sequence

$$H^{q-2}(Z, q-1) \xrightarrow{\alpha} H^q(X, q) \rightarrow H^q(U, q) \rightarrow H^{q-1}(Z, q-1) \xrightarrow{\beta} H^{q+1}(X, q)$$

where α and β are the Gysin maps. The assumption $q \neq \dim(X)$ implies $q-1 \neq \dim(Z)$ and the Lefschetz theory implies that α is surjective and β is injective so that $H^q(U, q) = 0$. If $q = 1$ the desired assertion follows from the exact sequence

$$0 \rightarrow H^1(U, 1) \rightarrow \bigoplus_{1 \leq j \leq s} H^0(Z_j, 0) \rightarrow H^2(X, 1),$$

where the last map is the Gysin map and the injectivity of the first map follows from the vanishing of $H^1(X, 1)$ by the assumption on X . Now assuming $\dim(X) > 1$, $s > 1$ and $q > 1$, we proceed by the double induction on $\dim(X)$ and s . Put

$$U' = X \setminus \bigcup_{j=2}^s Z_j, \quad W = U' - U = Z_1 \setminus (Z_1 \cap \bigcup_{j=2}^s Z_j).$$

We have the short exact sequence

$$H^q(U', q) \rightarrow H^q(U, q) \rightarrow H^{q-1}(W, q-1).$$

By induction $H^q(U', q) = H^q(U', q)_{triv}$ and $H^{q-1}(W, q-1) = H^{q-1}(W, q-1)_{triv}$ while it is easy to check that the residue map $H^q(U, q)_{triv} \rightarrow H^{q-1}(W, q-1)_{triv}$ is surjective. This proves the desired assertion for $H^q(U, q)$.

To show Lemma2.3(2) let $H^q(V, q)_{triv} \subset H^q(V, q)$ be defined as $H^q(U, q)_{triv}$ by taking $X = \mathbb{P}^n$. By definition we have $H^q(U, q)_{triv} = \text{Image}(H^q(V, q)_{triv} \rightarrow H^q(U, q))$. By Lemma2.3(3) $H^q(V, q)_{triv} = H^q(V, q)$ if $q \neq n$. This proves the desired assertion.

3. Hodge theoretic implication of generalized Jacobian rings

In this section we prove Theorem 0.4 by using the results in §1. Let the assumption be as in §2. We fix a non-singular affine algebraic variety S over k and the following schemes over S

$$(3-1) \quad \mathcal{Y} = \bigcup_{1 \leq j \leq s} \mathcal{Y}_j \hookrightarrow \mathbb{P}_S^n \hookleftarrow \mathcal{X} \hookleftarrow \mathcal{Z} = \bigcup_{1 \leq j \leq s} \mathcal{Z}_j, \quad \mathcal{V} = \mathbb{P}_S^n \setminus \mathcal{Y}, \quad \mathcal{U} = \mathcal{X} \setminus \mathcal{Z}$$

whose fibers are as (2-1). We assume that the fibers of $\mathcal{X}/S$ are smooth complete intersections of multi-degree $(d_1, \dots, d_r)$ and those of $\mathcal{Y}_j \subset \mathbb{P}_S^n$ for $1 \leq j \leq s$ are hypersurfaces of degree e_j . For integers p, q we introduce the following sheaf on S_{zar}

$$H^{p,q}(\mathcal{U}/S) = R^q f_* \Omega_{\mathcal{X}/S}^p(\log \mathcal{Z}),$$

where $f : \mathcal{X} \rightarrow S$ is the natural morphism and $\Omega_{\mathcal{X}/S}^p(\log \mathcal{Z}) = \bigwedge^p \Omega_{\mathcal{X}/S}^1(\log \mathcal{Z})$ with $\Omega_{\mathcal{X}/S}^1(\log \mathcal{Z})$, the sheaf of relative differentials on $\mathcal{X}$ over S with logarithmic poles along $\mathcal{Z}$. In case $s \geq 1$ the Lefschetz theory implies $H^{p,q}(\mathcal{U}/S) = 0$ if $p + q \neq n - r$. In case $s = 0$ it implies $H^{p,q}(\mathcal{X}/S)_{prim} = 0$ if $p + q \neq n - r$ where “*prim*” denotes the primitive part (cf. Theorem 1.3(1)). The results in §1 implies that under an appropriate numerical condition on d_i and e_j we can control the cohomology of the following Koszul complex

$$\Omega_S^{q-1} \otimes H^{a+2,b-2}(\mathcal{U}/S) \xrightarrow{\overline{\nabla}} \Omega_S^q \otimes H^{a+1,b-1}(\mathcal{U}/S) \xrightarrow{\overline{\nabla}} \Omega_S^{q+1} \otimes H^{a,b}(\mathcal{U}/S).$$

Here $\overline{\nabla}$ is induced by the Kodaira-Spencer map

$$(3-1) \quad \kappa_{(\mathcal{X}, \mathcal{Z})} : \Theta_S \rightarrow R^1 f_* T_{\mathcal{X}/S}(-\log \mathcal{Z}),$$

with $\Theta_S = \mathcal{H}om_{\mathcal{O}_S}(\Omega_S^1, \mathcal{O}_S)$ and $T_{\mathcal{X}/S}(-\log \mathcal{Z}) = \mathcal{H}om_{\mathcal{O}_{\mathcal{X}}}(\Omega_{\mathcal{X}/S}^1(\log \mathcal{Z}), \mathcal{O}_{\mathcal{X}})$, and the map

$$R^1 f_* T_{\mathcal{X}/S}(-\log \mathcal{Z}) \otimes R^{b-1} f_* \Omega_{\mathcal{X}/S}^{a+1}(\log \mathcal{Z}) \rightarrow R^b f_* \Omega_{\mathcal{X}/S}^a(\log \mathcal{Z})$$

induced by the cup product and $T_{\mathcal{X}/S}(-\log \mathcal{Z}) \otimes \Omega_{\mathcal{X}/S}^{a+1}(\log \mathcal{Z}) \rightarrow \Omega_{\mathcal{X}/S}^a(\log \mathcal{Z})$, the contraction. For the application to the Beilinson’s conjectures the kernel of the following map plays a crucial role

$$\overline{\nabla}^{p,q} : H^{p,q}(\mathcal{U}/S) \rightarrow \Omega_S^1 \otimes_{\mathcal{O}} H^{p-1,q+1}(\mathcal{U}/S).$$

In case $s = 0$ we let $\overline{\nabla}^{p,q}$ denote the primitive part of the above map:

$$H^{p,q}(\mathcal{X}/S)_{prim} \xrightarrow{\overline{\nabla}} \Omega_S^1 \otimes_{\mathcal{O}} H^{p-1,q+1}(\mathcal{X}/S)_{prim}.$$

Now we fix a k -rational point $0 \in S$ and let $U \subset X \supset Z$ denote the fibers of $\mathcal{U} \subset \mathcal{X} \supset \mathcal{Z}$. Taking the fibers of the above maps we get

$$\begin{aligned} \overline{\nabla}_0^{p,q} : H^{p,q}(U) &\rightarrow \Omega_{S,0}^1 \otimes H^{p-1,q+1}(U) \quad (s \geq 1), \\ \overline{\nabla}_0^{p,q} : H^{p,q}(X)_{prim} &\rightarrow \Omega_{S,0}^1 \otimes H^{p-1,q+1}(X)_{prim} \quad (s = 0) \end{aligned}$$

where $H^{p,q}(U) = H^q(X, \Omega_X^p(\log Z))$ and $H^{p,q}(X)_{\text{prim}} = H^q(X, \Omega_X^p)_{\text{prim}}$. Let $T_0(S)$ be the tangent space of S at 0 and fix a linear subspace $T \subset T_0(S)$. Via the canonical isomorphism $\Omega_{S,0}^1 \cong \text{Hom}_k(T_0(S), k)$, the above maps induce

$$\begin{aligned}\overline{\nabla}_{0,T}^{p,q} &: H^{p,q}(U) \rightarrow \text{Hom}_k(T, H^{p-1,q+1}(U)) \quad (s \geq 1), \\ \overline{\nabla}_{0,T}^{p,q} &: H^{p,q}(X)_{\text{prim}} \rightarrow \text{Hom}_k(T, H^{p-1,q+1}(X)_{\text{prim}}) \quad (s = 0)\end{aligned}$$

Now the key result is the following. See Definition 3.2 below for the definition of $c_S(\mathcal{X}, \mathcal{Z})_0$. Put

$$\mathbf{d} = \sum_{1 \leq i \leq r} d_i, \quad \delta_{\min} = \min\{d_i, e_j \mid 1 \leq j \leq s, 1 \leq i \leq r\}.$$

Theorem 3.1 *Assume $p + q = m := n - r \geq 1$. Let $c = \text{codim}_{T_0(S)}(T)$.*

- (1) *Assuming $1 \leq p \leq m - 1$ and $\delta_{\min}(p - 1) + \mathbf{d} \geq n + 1 + c_S(\mathcal{X}, \mathcal{Z})_0 + c$, $\text{Ker}(\overline{\nabla}_{0,T}^{p,q}) = 0$.*
- (2) *Assuming $\delta_{\min}(n - r - 1) + \mathbf{d} \geq n + 1 + c_S(\mathcal{X}, \mathcal{Z})_0 + c$, $\text{Ker}(\overline{\nabla}_{0,T}^{m,0}) = H^0(X, \Omega_X^m(\log Z))_{\text{triv}}$ (cf. Definition 2.1).*

Definition 3.2 *We define*

$$c_S(\mathcal{X}, \mathcal{Z})_0 = \dim_k(\text{Image}(\psi_{(X,Z)}) / \text{Image}(\psi_{(X,Z)}) \cap \text{Image}(\kappa_{(X,Z)})).$$

where $\kappa_{(X,Z)} : T_0(S) \rightarrow H^1(X, T_X(-\log Z))$ (resp. $\psi_{(X,Z)} : B_1(0) \rightarrow H^1(X, T_X(-\log Z))$) is the fiber at 0 of the map (3-2) (resp. the map in Theorem 1.3(2) for (X, Z)). We also define

$$c_S(\mathcal{X}, \mathcal{Z}) = \min_{t \in S} \{c_S(\mathcal{X}, \mathcal{Z})_t\}$$

where $c_S(\mathcal{X}, \mathcal{Z})_t$ is defined by the same way as $c_S(\mathcal{X}, \mathcal{Z})_0$ for the fibers of $\mathcal{U} \subset \mathcal{X} \supset \mathcal{Z}$ over $t \in S$.

Remark 3.3 *If $n - r \geq 2$ and X is not a K3 surface, $\psi_{(X,Z)}$ is surjective so that $c_S(\mathcal{X}, \mathcal{Z})_0 = \dim_k \text{Coker}(\kappa_{(X,Z)})$.*

Let $P = \mathbb{C}[z_0, \dots, z_n]$ be the homogeneous coordinate ring of $\mathbb{P}^n$ and let $P^d \subset P$ be the space of the homogeneous polynomials of degree d . The dual projective space

$$\mathbb{P}^\vee(P^d) = \mathbb{P}_k^{N(n,d)} \quad (N(n,d) = \binom{n+d}{d} - 1)$$

parametrizes hypersurfaces $Y \subset \mathbb{P}^n$ of degree d defined over k . Let

$$B \subset \prod_{1 \leq i \leq r} \mathbb{P}_k^{N(n,d_i)} \times \prod_{1 \leq j \leq s} \mathbb{P}_k^{N(n,e_j)}$$

be the Zariski open subset parametrizing such $((X_i)_{1 \leq i \leq r}, (Y_j)_{1 \leq j \leq s})$ that $X_1 + \dots + X_r + Y_1 + \dots + Y_s$ is a simple normal crossing divisor on $\mathbb{P}_k^n$. We consider the family

$$\mathcal{X}_B \hookrightarrow \mathcal{Z}_B = \bigcup_{1 \leq j \leq s} \mathcal{Z}_{B,j} \quad \text{over } B$$

whose fibers are $X \hookrightarrow Z = \bigcup_{1 \leq j \leq s} Z_j$ with $X = X_1 \cap \dots \cap X_r$ and $Z_j = X \cap Y_j$. We omit the proof of the following.

Lemma 3.4 *Let $E \subset B$ be a non-singular locally closed subvariety of codimension $c \geq 0$ and let $S \rightarrow E$ be a dominant map. Assume that the family $(\mathcal{X}, \mathcal{Z})/S$ is the pullback of $(\mathcal{X}_B, \mathcal{Z}_B)/B$ via $S \rightarrow B$. Then we have $c_S(\mathcal{X}, \mathcal{Z}) \leq c$.*

Now we prove Theorem3.1. We only show the second assertion and leave the first to the readers. The fact $H^0(X, \Omega_X^m(\log Z))_{\text{triv}} \subset \text{Ker}(\overline{\nabla}_{0,T}^{m,0})$ follows from the fact that it lies in the image of

$$H^0(\mathcal{X}, \Omega_{\mathcal{X}/k}^m(\log \mathcal{Z})) \rightarrow \Gamma(S, f_* \Omega_{\mathcal{X}/S}^m(\log \mathcal{Z})) \rightarrow H^0(X, \Omega_X^m(\log Z)),$$

where $\Omega_{\mathcal{X}/k}(\log \mathcal{Z})$ is the sheaf of differential forms of $\mathcal{X}$ over k with logarithmic poles along $\mathcal{Z}$. Thus we are reduced to show the injectivity of the induced map

$$H^0(X, \Omega_X^m(\log Z))/H^0(X, \Omega_X^m(\log Z))_{\text{triv}} \rightarrow \text{Hom}_k(T, H^1(X, \Omega_X^{m-1}(\log Z))).$$

By Theorem1.5 and Theorem1.6 in §1 this is reduced to show the surjectivity of

$$W \otimes B_{n-r-1}(\mathbf{d} - n - 1) \rightarrow B_{n-r}(\mathbf{d} - n - 1)$$

where $B_1(0) \supset W := \psi_{(X,Z)}^{-1}(\kappa_{(X,Z)}(T))$. By definition W is of codimension $\leq c + c_S(\mathcal{X}, \mathcal{Z})_0$ in $B_1(0)$. Hence the desired assertion follows from Theorem1.8(i) in §1.

Having Theorem3.1, Theorem0.4 is proven by a standard method in Hodge theory (cf. [G1, p.75]), which we recall in what follows. Let the assumption be as in Theorem0.4. Fix $0 \in E \subset S_{NL}$ and let $U \subset X \supset Z$ be as before. Choosing a simply connected neighbourhood Δ of 0 in $S(\mathbb{C})$ we have the natural identification for $\forall t \in \Delta$ via flat translation with respect to the Gauss-Maninn connection:

$$H^m(U, \mathbb{Q}(m)) \xrightarrow{\cong} H^m(U_t, \mathbb{Q}(m)); \lambda \rightarrow \lambda_t$$

where U_t is the fiber of $\mathcal{U}$ over $t \in S$. By definition of S_{NL} there exists $\lambda \in F^0 H^m(U, \mathbb{Q}(m))$ not contained in $H^m(U, \mathbb{Q}(m))_{\text{triv}}$ such that $\lambda_t \in F^m H^m(U_t, \mathbb{C})$ for $\forall t \in E \cap \Delta$. By noting Remark2.2 it implies that $\lambda \in \text{Ker}(\overline{\nabla}_{0,T}^{m,0}) \neq H^0(X, \Omega_X^m(\log Z))_{\text{triv}}$, where $T \subset T_0(S)$ is the tangent space of E . Hence we have

$$\text{codim}_S(E) \geq \text{codim}_{T_0(S)}(T) \geq \delta_{\min}(n - r - 1) + \sum_{1 \leq i \leq r} d_i - n - c_S(\mathcal{X}, \mathcal{Z})$$

where the second inequality follows from Theorem3.1. This completes the proof of Theorem0.4.

4. Case of plane curves

One may naturally asks if the estimate in Theorem0.4 is optimal. In this section we answer the question in case $n = 2$ and $r = 1$, namely in case the fibers of $\mathcal{X}/S$ are plane curves. We work over $\mathbb{C}$. Let $P = \mathbb{C}[z_0, z_1, z_2]$ be the homogeneous coordinate ring of $\mathbb{P}^2$ and let $P^d \subset P$ be the space of the homogeneous polynomials of degree d . Let

$$S \subset P^d/\mathbb{C}^* \times \prod_{1 \leq j \leq s} P^{e_j}/\mathbb{C}^* = \mathbb{P}^{N(2,d)} \times \prod_{1 \leq j \leq s} \mathbb{P}^{N(2,e_j)} \quad (N(2,u) = \binom{u+2}{2} - 1)$$

be the moduli space of such sets of plane curves $(X, (Y_j)_{1 \leq j \leq s})$ that $X \cup Y_1 \cup \dots \cup Y_s$ is a simple normal crossing divisor on $\mathbb{P}^2$. Let $\mathcal{X} \hookrightarrow \mathbb{P}_S^n \hookleftarrow \mathcal{Y}_j$ be the universal families over S and put $\mathcal{Z} = \mathcal{X} \cap \bigcup_{1 \leq j \leq s} \mathcal{Y}_j$ and $\mathcal{U} = \mathcal{X} \setminus \mathcal{Z}$. Let $U_t \subset X_t \supset Z_t$ denote the fibers of $\mathcal{U} \subset \mathcal{X} \supset \mathcal{Z}$ over $t \in S$. We are interested in the Noether-Lefschetz locus for Beilinson-Hodge cycles on U_t ;

$$S_{NL} = \{t \in S \mid F^0 H^1(U_t, \mathbb{Q}(1)) \neq H^1(U_t, \mathbb{Q}(1))\}.$$

Noting that $\text{reg}_{U_t}^1$ is known to be surjective and that $\text{Ker}(\text{reg}_{U_t}^1) = \mathbb{C}^* \otimes \mathbb{Q} \subset CH^1(U_t, 1) \otimes \mathbb{Q}$, we have

$$S_{NL} = \{t \in S \mid CH^1(U_t, 1) \otimes \mathbb{Q} \neq CH^1(U_t, 1)_{\text{triv}}\}.$$

Theorem 4.1 *Let $E \subset S_{NL}$ be an irreducible component. Then $\text{codim}_S(E) \geq d - 2$. Moreover there exists E for which the equality holds.*

Proof. The inequality follows from Theorem 0.4 together with the fact $c_S(\mathcal{X}, \mathcal{Z}) = 0$, which follows from Lemma 3.4. As for the second assertion we consider the subset $\Sigma \subset S$ of those $t \in S$ that satisfy the condition: there exists the unique point $x_t \in Z_t$ such that $X_t \cap H_t = \{x_t\}$ set-theoretically, where H_t is the tangent line of X_t at x_t . It is easy to check that Σ is locally closed algebraic subset of S . Thus it suffices to show the following.

Lemma 4.2 $\Sigma \subset S_{NL}$, Σ is equidimensional, and $\text{codim}_S(\Sigma) = d - 2$.

Proof. Fix any $t \in \Sigma$ and let Y_j and U denote the fibers of $\mathcal{Y}_j$ and $\mathcal{U}$ over t for simplicity. Choose $L \in P^1$ defining $H_t \subset \mathbb{P}^2$ and $G_j \in P^{e_j}$ defining Y_j which contains x_t . Then it is easy to see that $(G_j/L^{e_j})|_U \in CH^1(U, 1)$ while it is not contained in $CH^1(U, 1)_{\text{triv}}$. This proves the first assertion of Lemma 4.2. In order to show the remaining assertions, let M be the space of pairs (x, H) of a point $x \in \mathbb{P}^2$ and a line $H \subset \mathbb{P}^2$ passing through x . Define the morphism $\pi : \Sigma \rightarrow M$ by $\pi(t) = (x_t, H_t)$. Considering the action of $\text{PGL}_3(\mathbb{C})$, the group of linear transformations on $\mathbb{P}^2$, one easily see that π is surjective. Thus it suffices to show the following.

Lemma 4.3 *Let $\Sigma_0 \subset \Sigma$ be the fiber of π over $(0, H_0)$ with $0 := (0 : 0 : 1)$ and $H_0 \subset \mathbb{P}^2$ defined by $z_0 \in P^1$. Then Σ_0 is equidimensional and $\text{codim}_S(\Sigma) = d - 2 + \dim(M) = d + 1$.*

Proof. Let $T \subset S' := \prod_{1 \leq j \leq s} \mathbb{P}_k^{N(2, e_j)}$ be the closed subset parametrizing such $(Y_j)_{1 \leq j \leq s}$ that $0 \in Y_j$ for some j . Clearly T is equidimensional and of $\text{codim}_{S'}(T) = 1$. By definition one sees that X_t for $t \in E_0$ is defined by an equation of the form $z_0 A + z_1^d$ with $A \in P^{d-1}$. This implies that E_0 is identified with a non empty open subset of $P^{d-1} \times T$, where P^{d-1} is viewed as an affine space over $\mathbb{C}$. This shows that E_0 is equidimensional and that $\text{codim}_S(E_0) = \binom{d+2}{2} - 1 - \binom{d+1}{2} + \text{codim}_{S'}(T) = d + 1$ as desired.

Remark 4.4 *It is interesting to ask if the components E of Σ are the only ones satisfying $\text{codim}_S(E) = d - 2$. It is possible to give a positive answer to this question in case $s = 2$ and $e_1 = e_2 = 1$ by using the method in [Gr5] and [V]. The general case remains open.*

5. Beilinson's Tate conjecture

In this section we show Theorem0.5. Let the assumption be as in the beginning of §3. Write $m = n - r$ and $\mathbf{d} = \sum_{1 \leq i \leq r} d_i$. Theorem0.5(1) is an immediate consequence of Theorem5.1 and Remark5.2 below.

Theorem 5.1 *Assume that k is finitely generated over $\mathbb{Q}$ and that $\mathbf{d} \geq n + 1 + c_S(\mathcal{X}, \mathcal{Z})$. Let $\bar{k}$ be an algebraic closure of k and put $\bar{S} = S \times_k \bar{k}$. Let $\bar{\eta}$ be a geometric generic point of $\bar{S}$ and let $U_{\bar{\eta}} \subset X_{\bar{\eta}}$ be the fibers of $\mathcal{U} \subset \mathcal{X}$ over $\bar{\eta}$.*

- (1) *Assuming $s \geq 1$, $H_{et}^m(U_{\bar{\eta}}, \mathbb{Q}_\ell(m))^{\pi_1(\bar{S}, \bar{\eta})} = H_{et}^m(U_{\bar{\eta}}, \mathbb{Q}_\ell(m))_{triv}$.*
- (2) *$H_{et}^m(X_{\bar{\eta}}, \mathbb{Q}_\ell(m))_{prim}^{\pi_1(\bar{S}, \bar{\eta})} = 0$.*

Remark 5.2 *Let $\tilde{S} \rightarrow S$ be a dominant morphism and $\tilde{\mathcal{X}} \supset \tilde{\mathcal{Z}}$ be the base change. Then $c_S(\mathcal{X}, \mathcal{Z}) = c_{\tilde{S}}(\tilde{\mathcal{X}}, \tilde{\mathcal{Z}})$. Hence Theorem5.1 holds after replacing $\mathcal{X} \supset \mathcal{Z}$ by the base change. In particular Theorem5.1 holds if one replaces $\pi_1(\bar{S}, \bar{\eta})$ by any open subgroup of finite index.*

Theorem 5.3 *Assume that $k = \mathbb{C}$ and $\mathbf{d} \geq n + 1 + c_S(\mathcal{X}, \mathcal{Z})$. Let $U \subset X$ be the fibers of $\mathcal{U} \subset \mathcal{X}$ over a fixed base point $0 \in S(\mathbb{C})$.*

- (1) *Assuming $s \geq 1$, $H^m(U, \mathbb{Q}(m))^{\pi_1(S, 0)} = H^m(U, \mathbb{Q}(m))_{triv}$.*
- (2) *$H^m(X, \mathbb{Q}(m))_{prim}^{\pi_1(S, 0)} = 0$.*

First we deduce Theorem5.1 from Theorem5.3. By the Lefschetz principle we may assume that k is a subfield of $\mathbb{C}$. We fix an embedding $\bar{k}(\bar{\eta}) \hookrightarrow \mathbb{C}$ and let $0 \in \bar{S}(\mathbb{C})$ be the corresponding $\mathbb{C}$ -valued point of $\bar{S}$. Write $S_{\mathbb{C}} = S \otimes_k \mathbb{C}$ and put $U_{\mathbb{C}} = \mathcal{X} \times_S \text{Spec}(\mathbb{C})$ via $\text{Spec}(\mathbb{C}) \xrightarrow{0} \bar{S} \rightarrow S$. We have the comparison isomorphisms ([SGA4, XVI Theorem4.1])

$$H^m(U_{\mathbb{C}}, \mathbb{Q}) \otimes \mathbb{Q}_\ell \xrightarrow{\cong} H_{et}^m(U_{\mathbb{C}}, \mathbb{Q}_\ell) \xrightarrow{\cong} H_{et}^m(U_{\bar{\eta}}, \mathbb{Q}_\ell)$$

that are equivariant with respect to the maps of topological and algebraic fundamental groups:

$$\pi_1^{top}(S_{\mathbb{C}}, 0) \rightarrow \pi_1^{alg}(S_{\mathbb{C}}, 0) \rightarrow \pi_1^{alg}(\bar{S}, \bar{\eta}).$$

The desired assertion follows at once from this.

In order to show Theorem5.3 we need a preliminary. We assume $k = \mathbb{C}$. Let S_{an} be the analytic site on $S(\mathbb{C})$. For a coherent sheaf $\mathcal{F}$ on S_{zar} let $\mathcal{F}^{an}$ be the associated analytic sheaf on S_{an} . We introduce local systems on S_{an}

$$H_{\mathbb{Q}}^q(\mathcal{U}/S)(p) = R^q g_* \mathbb{Q}(p) \quad \text{and} \quad H_{\mathbb{C}}^q(\mathcal{U}/S) = R^q g_* \mathbb{C},$$

where $g : \mathcal{U} \rightarrow S$ is the natural morphism. Let $H_{\mathbb{O}}^q(\mathcal{U}/S)$ be the sheaf of holomorphic sections of $H_{\mathbb{C}}^q(\mathcal{U}/S)$ and let $F^p H_{\mathbb{O}}^q(\mathcal{U}/S) \subset H_{\mathbb{O}}^q(\mathcal{U}/S)$ be the holomorphic subbundle given by the Hodge filtration on the cohomology of fibers of $\mathcal{U}/S$. We have the analytic Gauss-Manin connection

$$\nabla : H_{\mathbb{O}}^q(\mathcal{U}/S) \rightarrow \Omega_{S_{an}}^1 \otimes H_{\mathbb{O}}^q(\mathcal{U}/S) \quad \text{with} \quad \text{Ker}(\nabla) = H_{\mathbb{O}}^q(\mathcal{U}/S)$$

that satisfies $\nabla(F^p H_{\mathbb{O}}^q(\mathcal{U}/S)) \subset \Omega_{S_{an}}^1 \otimes F^{p-1} H_{\mathbb{O}}^q(\mathcal{U}/S)$. The induced map

$$F^p H_{\mathbb{O}}^{p+q}(\mathcal{U}/S)/F^{p+1} \rightarrow \Omega_{S_{an}}^1 \otimes F^{p-1} H_{\mathbb{O}}^{p+q}(\mathcal{U}/S)/F^p$$

is identified with $(\overline{\nabla}^{p,q})^{an}$ via the identification $F^p H_{\mathbb{O}}^{p+q}(\mathcal{U}/S)/F^{p+1} = H^{p,q}(\mathcal{U}/S)^{an}$ where

$$\overline{\nabla}^{p,q} : H^{p,q}(\mathcal{U}/S) \rightarrow \Omega_S^1 \otimes_{\mathcal{O}} H^{p-1,q+1}(\mathcal{U}/S)$$

is defined in §3. Therefore Theorem3.1 implies the following.

Theorem 5.4 *Let the assumption be as in Theorem5.3.*

(1) *If $s \geq 1$, $F^1 H_{\mathbb{O}}^m(\mathcal{U}/S) \cap H_{\mathbb{C}}^m(\mathcal{U}/S)$ is generated over $\mathbb{C}$ by the image of*

$$H^m(U, \mathbb{Q}(m))_{triv} \hookrightarrow H^m(U, \mathbb{Q}(m))^{\pi_1(S,0)} \simeq \Gamma(S_{an}, H_{\mathbb{Q}}^m(\mathcal{U}/S)(m)) \hookrightarrow H_{\mathbb{C}}^m(\mathcal{U}/S).$$

(2) *If $s = 0$, $F^1 H_{\mathbb{O}}^m(\mathcal{X}/S) \cap H_{\mathbb{C}}^m(\mathcal{X}/S)_{prim} = 0$, where $H_{\mathbb{C}}^q(\mathcal{X}/S)_{prim}$ is the primitive part of $H_{\mathbb{C}}^q(\mathcal{X}/S)$.*

Now we start the proof of Theorem5.3. First we show Theorem5.3(2). Write $H = H^m(X, \mathbb{Q})_{prim}^{\pi_1(S,0)}$. We have the natural isomorphism

$$H \otimes \mathbb{C} \simeq \Gamma(S_{an}, H_{\mathbb{C}}^m(\mathcal{X}/S)_{prim}).$$

By [D1] H is a sub-Hodge structure of $H^m(X, \mathbb{Q})$ so that we have the Hodge decomposition

$$H \otimes \mathbb{C} = \bigoplus_{p+q=m} H^{p,q} \quad \text{with} \quad H^{p,q} \subset \Gamma(S_{an}, F^p H_{\mathbb{O}}^m(\mathcal{X}/S) \cap H_{\mathbb{C}}^m(\mathcal{X}/S)_{prim}).$$

By Theorem5.4 this implies $H^{p,q} = 0$ for $p \geq 1$ which implies $H = 0$ by the Hodge symmetry.

Next we show Theorem5.3(1). By Lemma5.5 below we have

$$H^m(U, \mathbb{Q}(m))^{\pi_1(S,0)} \simeq \Gamma(S_{an}, H_{\mathbb{Q}}^m(\mathcal{U}/S)(m)) \subset \Gamma(S_{an}, H_{\mathbb{C}}^m(\mathcal{U}/S) \cap F^1 H_{\mathbb{O}}^m(\mathcal{U}/S)).$$

Hence the assertion follows from Theorem5.4(1).

Lemma 5.5 *Let the assumption be as in Theorem5.3. Then $H := H^m(U, \mathbb{Q})^{\pi_1(S, 0)}$ is a submixed Hodge structure of $H^m(U, \mathbb{Q})$ and $F^1 H_{\mathbb{C}} = H_{\mathbb{C}}$ where $H_{\mathbb{C}} = H \otimes \mathbb{C}$.*

Proof. The fact that H is a submixed Hodge structure follows from the theory of mixed Hodge modules [SaM]. We show the second assertion. We recall that the graded subquotients of the weight filtration of $H^m(U, \mathbb{Q})$ is given by

$$Gr_{m+p}^W H^m(U, \mathbb{Q}) = \bigoplus_{1 \leq j_1 < \dots < j_p \leq s} H^{m-p}(Z_{j_1} \cap \dots \cap Z_{j_p}, \mathbb{Q}(-p))_{\text{prim}}. \quad (0 \leq p \leq m)$$

Therefore it suffices to note that $H^m(X, \mathbb{Q})_{\text{prim}}^{\pi_1(S, 0)} = 0$ by Theorem5.3(2).

Now we show Theorem0.5(2). It will follow from Theorem5.6 below by using the theory of Hilbert set (cf. [La]).

Theorem 5.6 *Under the assumption of Theorem0.5(2), there exists an irreducible variety $\tilde{S}$ over k with a finite etale covering $\phi : \tilde{S} \rightarrow S$ for which the following holds: Let $\tilde{\pi} = \pi \circ \phi$ and let $H \subset \mathbb{P}^N(k)$ be the subset of such points that $\tilde{\pi}^{-1}(y)$ is irreducible. For $\forall x \in S$ such that $\pi(x) \in H$ and for any subgroup $G \subset \text{Gal}(k(\bar{x})/k(x))$ of finite index we have*

$$H_{\text{et}}^m(U_{\bar{x}}, \mathbb{Q}_{\ell}(m))^G = H_{\text{et}}^m(U_{\bar{x}}, \mathbb{Q}_{\ell}(m))_{\text{triv}},$$

where $\bar{x}$ is a geometric point over x and $U_{\bar{x}}$ is the fiber of $\mathcal{U}/S$ over $\bar{x}$.

Proof. (cf. [T] and [BE]) By choosing a k -rational point 0 of S , we get the decomposition

$$\pi_1(S, \bar{\eta}) = \pi_1(\bar{S}, \bar{\eta}) \times \text{Gal}(\bar{k}/k),$$

where the notation is as in Theorem5.1 and $\pi_1(\bar{S}, \bar{\eta})$ is identified with the quotient of $\pi_1(S, \bar{\eta})$ which classifies the finite etale coverings of S that completely decompose over 0 . Let

$$\Gamma = \text{Image}(\pi_1(\bar{S}, \bar{\eta}) \rightarrow \text{GL}_{\mathbb{Q}_{\ell}}(H_{\text{et}}^m(U_{\bar{\eta}}, \mathbb{Q}_{\ell}(m))).$$

From the fact that Γ contains an ℓ -adic Lie group as a subgroup of finite index, we have the following fact (cf. [T]): There exists a subgroup $\Gamma' \subset \Gamma$ of finite index such that a continuous homomorphism of pro-finite group $P \rightarrow \Gamma$ is surjective if and only if $P \rightarrow \Gamma \rightarrow \Gamma/\Gamma'$ is surjective as a map of sets. Let $\phi : \tilde{S} \rightarrow S$ be a finite etale covering that corresponds to the inverse image of Γ' in $\pi_1(\bar{S}, \bar{\eta})$ and let H be defined as in Theorem5.6. Fix $x \in S$ with $\pi(x) \in H$ and let $\bar{x}$ be a geometric point of x . By choosing a “path” $\bar{x} \rightarrow \bar{\eta}$, we get the map $\text{Gal}(k(\bar{x})/k(x)) = \pi_1(x, \bar{x}) \xrightarrow{\iota} \pi_1(S, \bar{\eta})$. By the definition of H the composite of ι with $\pi_1(S, \bar{\eta}) \rightarrow \Gamma \rightarrow \Gamma/\Gamma'$ is surjective so that $\text{Gal}(k(\bar{x})/k(x))$ surjects onto Γ by the above fact. It implies that for any subgroup $G \subset \text{Gal}(k(\bar{x})/k(x))$ of finite index, its image π in Γ is of finite index and hence that we have

$$H_{\text{et}}^m(U_{\bar{x}}, \mathbb{Q}_{\ell}(m))^G \xrightarrow{\cong} H_{\text{et}}^m(U_{\bar{\eta}}, \mathbb{Q}_{\ell}(m))^{\pi} = H_{\text{et}}^m(U_{\bar{\eta}}, \mathbb{Q}_{\ell}(m))_{\text{triv}}$$

where the last equality follows from Theorem5.1 and Remark5.2. This completes the proof of Theorem5.6.

Now Theorem 0.4(2) is a consequence of Theorem 5.6 and the following result (cf. [La]):

Lemma 5.7 *Let k be a number field and let V be an irreducible variety over k . Let $\pi : V \rightarrow \mathbb{P}_k^N$ be an étale morphism and let $H \subset \mathbb{P}_k^N(k)$ be the subset of such points x that $\pi^{-1}(x)$ is irreducible. Let Σ be any finite set of primes of k and let k_v be the completion of k at v . Then the image of H in $\prod_{v \in \Sigma} \mathbb{P}_k^N(k_v)$ is dense.*

6. Implication on injectivity of regulator maps for K_1 of surfaces

Let X be a projective smooth surface over a field k . Let $U \subset X$ be the complement of a simple normal crossing divisor $Z \subset X$. In this section we discuss an implication of the surjectivity of reg_U^2 and $\text{reg}_{\text{ét}, U}^2$ on the regulator maps for $CH^2(X, 1)$. Recall that $CH^2(X, 1)$ is by definition the cohomology of the following complex

$$K_2(k(X)) \xrightarrow{\partial_{\text{tame}}} \bigoplus_{C \subset X} k(C)^* \xrightarrow{\partial_{\text{div}}} \bigoplus_{x \in X} \mathbb{Z},$$

where the sum on the middle term ranges over all irreducible curves on X and that on the right hand side over all closed points of X . The map ∂_{tame} is the so-called tame symbol and ∂_{div} is the sum of divisors of rational functions on curves. Thus an element of $\text{Ker}(\partial_{\text{div}})$ is given by a finite sum $\sum_i (C_i, f_i)$, where f_i is a non-zero rational function on an irreducible curve $C_i \subset X$ such that $\sum_i \text{div}(f_i) = 0$ on X . We recall that $K_2(k(X))$ is generated as an abelian groups by symbols $\{f, g\}$ for non-zero rational functions f, g on X and that

$$\partial_{\text{tame}}(\{f, g\}) = ((f)_0, g) + ((f)_\infty, 1/g) + ((g)_0, 1/f) + ((g)_\infty, f),$$

where $(f)_0$ (resp. $(f)_\infty$) is the zero (resp. pole) divisor of f . An important tool to study $CH^2(X, 1)$ is the regulator map: In case $k = \mathbb{C}$ it is given by

$$\text{reg}_{D, X} : CH^2(X, 1) \rightarrow H_D^3(X, \mathbb{Z}(2)),$$

where the group on the right hand side is the Deligne cohomology group (cf. [EV] and [J2]). Assuming the first Betti number $b_1(X) = 0$, we have the following explicit description of $\text{reg}_{D, X}$. Take $\alpha = \sum_i (C_i, f_i) \in \text{Ker}(\partial_{\text{div}})$. Under the isomorphism (cf. [EV, 2.10])

$$H_D^3(X, \mathbb{Z}(2)) \simeq \frac{H^2(X, \mathbb{C})}{H^2(X, \mathbb{Z}(2)) + F^2 H^2(X, \mathbb{C})} \simeq \frac{F^1 H^2(X, \mathbb{C})^*}{H_2(X, \mathbb{Z})},$$

$\text{reg}_{D, X}(\alpha)$ is identified with a linear function on complex valued C^∞ -forms ω and we have

$$\text{reg}_{D, X}(\alpha)(\omega) = \frac{1}{2\pi\sqrt{-1}} \sum_i \int_{C_i - \gamma_i} \log(f_i) \omega + \int_\Gamma \omega,$$

where $\gamma_i := f_i^{-1}(\gamma_0)$ with γ_0 , a path on $\mathbb{P}_\mathbb{C}^1$ connecting 0 with ∞ and Γ is a real piecewise smooth 2-chain on X such that $\partial\Gamma = \cup_i \gamma_i$ which exists due to the assumption $\alpha \in \text{Ker}(\partial_{\text{div}})$ and $b_1(X) = 0$.

In case k is finitely generated over $\mathbb{Q}$ we have the regulator map

$$\text{reg}_{\text{cont}, X} : CH^2(X, 1) \otimes \mathbb{Z}_\ell \rightarrow H_{\text{cont}}^3(X, \mathbb{Z}_\ell(2)),$$

where H_{cont}^i denotes the continuous étale cohomology of X (cf. [J3]).

Now let $Z \subset X \supset U$ be as in the begining of this section and write $Z = \cup_{1 \leq i \leq r} Z_i$ with Z_i , smooth irreducible curves intersecting transversally with each other. We consider the following subgroup of $\text{Ker}(\partial_{\text{div}})$

$$CH^1(Z, 1) = \text{Ker}\left(\bigoplus_{1 \leq i \leq r} k(Z_i)^* \xrightarrow{\partial_{\text{div}}} \bigoplus_{x \in Z} \mathbb{Z}\right).$$

By the localization theory for higher Chow group we have the exact sequence

$$CH^2(U, 2) \xrightarrow{\partial_Z} CH^1(Z, 1) \xrightarrow{\iota} CH^2(X, 1)$$

where the first map coincides up to sign with the composite of the natural map $CH^2(U, 2) \rightarrow K_2(k(X))$ and ∂_{tame} .

Theorem 6.1 (1) *Assume $k = \mathbb{C}$ and that there exists a subspace $\Delta \subset CH^2(U, 2) \otimes \mathbb{Q}$ such that the restriction of reg_U^2 on Δ is surjective. Let $\alpha \in CH^1(Z, 1)$ and assume $\text{reg}_{D,X}(\iota(\alpha)) = 0$. Then $\alpha \in \partial_Z(\Delta)$ in $CH^1(Z, 1) \otimes \mathbb{Q}$. In particular $\iota(\alpha) = 0 \in CH^2(X, 1) \otimes \mathbb{Q}$.*

(2) *Assume that k is finitely generated over $\mathbb{Q}$. The analogous fact holds for $\text{reg}_{\text{et},U}^2$ and $\text{reg}_{\text{cont},X}$.*

The following result is a direct consequence of Theorem 6.1 and Definition 0.3. Let $Z \subset X$ be as (0-4) in the introduction and let $Z_t \subset X_t$ be the fibers over $t \in S$.

Corollary 6.2 *Let the assumption be as in Theorem 0.4. For any $t \in S \setminus S_{NL}$, reg_{D,X_t} restricted on the image of $CH^1(Z_t, 1)$ is injective modulo torsion.*

Remark 6.3 *In the forthcoming paper [AS2] we study the Noether-Lefschetz locus in case (0-4) is the universal family of smooth surfaces of degree d and three hyperplanes in $\mathbb{P}^3$ intersecting transversally. In this case it is shown that there exist $t \in S \setminus S_{NL}$ such that the image of $CH^1(Z_t, 1)$ in $CH^2(X_t, 1)$ is non-torsion if d is large enough. Thus Corollary 6.2 has indeed non-trivial implication on the injectivity of the regulator map.*

Now we start the proof of Theorem 6.1. The idea of the following proof is taken from [J1, 9.8]. We only treat the first assertion. The second is proven by the same way. We have the commutative diagram (cf. [Bl] and [J2, 3.3 and 1.15]).

$$\begin{array}{ccccccc} CH^2(X, 2) & \rightarrow & CH^2(U, 2) & \rightarrow & CH^1(Z, 1) & \rightarrow & CH^2(X, 1) \\ \downarrow ch_{D,X}^{2,2} & & \downarrow ch_{D,U}^{2,2} & & \downarrow ch_{D,Z}^{1,1} & & \downarrow ch_{D,X}^{2,1} \\ H_D^2(X, \mathbb{Z}(2)) & \xrightarrow{\iota} & H_D^2(U, \mathbb{Z}(2)) & \rightarrow & H_{D,Z}^3(X, \mathbb{Z}(2)) & \rightarrow & H_D^3(X, \mathbb{Z}(2)) \end{array}$$

Here the horizontal sequences are the localization sequences for higher Chow groups and Deligne cohomology groups and they are exact. The vertical maps are regulator maps. By a simple diagram chasing it suffices to show that $ch_{D,Z}^{1,1}$ is injective and $ch_{D,U}^{2,2}(\Delta) + \text{Image}(\iota)$ spans $H_D^2(U, \mathbb{Q}(2))$. In order to show the first assertion we note the commutative diagram

$$\begin{array}{ccccccc} 0 \rightarrow & \bigoplus_{1 \leq i \leq r} CH^1(Z_i, 1) & \rightarrow & CH^1(Z, 1) & \rightarrow & \bigoplus_{x \in W} \mathbb{Z} \\ & \downarrow ch_{D,Z_i}^{1,1} & & \downarrow ch_{D,Z}^{1,1} & & \simeq \downarrow ch_{D,x}^{0,0} \\ 0 \rightarrow & \bigoplus_{1 \leq i \leq r} H_D^1(Z_i, \mathbb{Z}(1)) & \rightarrow & H_{D,Z}^3(X, \mathbb{Z}(2)) & \rightarrow & \bigoplus_{x \in W} H_D^0(x, \mathbb{Z}(0)) \\ & \downarrow \simeq & & \downarrow = & & \downarrow \simeq \\ & \bigoplus_{1 \leq i \leq r} H_{D,Z_i}^3(X, \mathbb{Z}(2)) & \rightarrow & H_{D,Z}^3(X, \mathbb{Z}(2)) & \rightarrow & \bigoplus_{x \in W} H_{D,x}^4(X, \mathbb{Z}(2)) \end{array}$$

where $W = \coprod_{1 \leq i \neq j \leq r} Z_i \cap Z_j$. The horizontal sequences come from the Mayer-Vietoris spectral sequence and they are exact. Thus the desired assertion follows from the fact that $ch_{D, Z_i}^{1,1}$ is an isomorphism (cf. [J2, 3.2]). To show the second assertion we recall the exact sequence (cf. [EV, 2.10])

$$0 \rightarrow H^1(U, \mathbb{C})/H^1(U, \mathbb{Z}(2)) \xrightarrow{\gamma} H_D^2(U, \mathbb{Z}(2)) \xrightarrow{\beta} H^2(U, \mathbb{Z}(2)) \cap F^2 H^2(U, \mathbb{C}) \rightarrow 0$$

and the same sequence with U replaced by X . We have $ch_{B,U}^{2,2} = \beta \cdot ch_{D,U}^{2,2}$ and the commutative diagram

$$\begin{array}{ccc} \mathbb{C}^* \otimes CH^1(U, 1) & \rightarrow & CH^1(U, 1) \otimes CH^1(U, 1) \\ \downarrow \log \otimes ch_{B,U}^{1,1} & & \downarrow \text{product} \\ \mathbb{C}/\mathbb{Z}(1) \otimes H^1(U, \mathbb{Z}(1)) & & CH^2(U, 2) \\ \downarrow \simeq & & \downarrow ch_{D,U}^{2,2} \\ H^1(U, \mathbb{C})/H^1(U, \mathbb{Z}(2)) & \xrightarrow{\gamma} & H_D^2(U, \mathbb{Z}(2)) \end{array}$$

Therefore the desired assertion follows from the surjectivity of

$$CH^1(U, 1) \xrightarrow{ch_{B,U}^{1,1}} H^1(U, \mathbb{Z}(1))/H^1(X, \mathbb{Z}(1))$$

which is easily seen.

References

- [AS1] M. Asakura and S. Saito, *Generalized Jacobian rigns for open complete intersections*, preprint
- [AS2] M. Asakura and S. Saito, *Noether-Lefschetz problem for Beilinson-Hodge cycles on open surfaces*, preprint
- [BE] S. Bloch and H. Esnault, *The coniveau filtration and non-divisibility for algebraic cycles*, Math. Ann. **304** (1996), 303–314
- [Bei] A. Beilinson, *Notes on absolute Hodge cohomology; Applications of Algebraic K-theory to Algebraic Geometry and Number theory*, Contemp. Math. **55** (1986), 35–68
- [Bl] S. Bloch, *Algebraic cycles and higher K-theory*, Advances in Math., **61** (1986), 267–304
- [CDK] E. Cattani, P. Deligne, A. Kaplan, *On the locus of Hodge classes*, Journal of the AMS, **8** (1995), 483–506
- [CGGH] J. Carlson, M. Green, P. Griffiths, J. Harris, *Infinitesimal variations of Hodge structure (I)*, Compositio Math., **50** (1983), 109–205
- [D1] P. Deligne, *Théorie de Hodge II*, Publ. Math. IHES, **40** (1972), 5–57
- [D2] P. Deligne *Le Théorème de Noether*, Lecture Notes in Math., **340** (1973), 328–340

- [Do] R. Donagi *Generic Torelli for projective hypersurfaces*, Compositio.Math., **50** (1983), 325-353
- [DG] R. Donagi and M. Green *A new proof of the symmetrizer lemma and a stronger weak Torelli theorem for projective hypersurfaces*, J.Diff.Geom., **20** (1984), 459-461
- [EV] H. Esnault and E. Viehweg, *Deligne-Beilinson cohomology*, in: Beilinson's Conjectures on Special Values of L -functions (M. Rapoport, N. Schappacher and P. Schneider, ed), Perspectives in Math., **4**, Academic Press
- [G1] M. Green *Infinitesimal methods in Hodge theory*, Lecture Notes in Math., **1594** (1993), 1-92 Springer-Verlag
- [G2] M. Green *The period map for hypersurface sections of high degree on an arbitrary variety*, Compositio Math., **55** (1984), 135-156
- [G3] M. Green *Koszul cohomology and Geometry*, Lectures on Riemann surfaces, Cornalba, Gomez-Mont, and Verjovsky (edotors), 177-200, ICTP, Trieste, Italy.
- [G4] M. Green, *A new proof of the explicit Noether-Lefschetz theorem*, J. Differential Geometry, **27** (1988), 155-159
- [G5] M. Green, *Components of maximal dimension in the Noether-Lefschetz locus*, J. Differential Geometry, **29** (1989), 295-302
- [Gri] P. Griffiths *Periods of certain rational integrals:I and II*, Ann. of Math., **90** (1969), 460-541
- [J1] U.Jannsen, *Mixed motives and Algebraic K-Theory*, Lecture Notes in Math., **1400** (1980), Springer-Verlag
- [J2] U.Jannsen, *Deligne homology, Hodge-D-conjecture, and Motives*, in: Beilinson's Conjectures on Special Values of L -functions (M. Rapoport, N. Schappacher and P. Schneider, ed), Perspectives in Math., **4**, Academic Press
- [J3] U.Jannsen, *Continuous etale cohomology*, Math. Annal., **280** (1987), 207-245
- [K] K.Konno, *On the variational Torelli problem for complete intersections*, Compositio Math., **78** (1991), 271-296
- [La] S. Lang *Sur les séries L d'une variété algébrique*, Bull. Soc. Math. France, **84** (1956), 385-407
- [Mu] D. Mumford *Rational equivalence of 0-cycles on surfaces*, J. Math. Kyoto Univ., **9** (1969), 195-204
- [Na] J. Nagel *The Abel-jacobi map for complete intersections*, Indag. Mathem., **8** (1997), 95-113
- [N] M. V. Nori, *Algebraic cycles and Hodge theoretic connectivity*, Invent. of Math., **111** (1993), 349-373

- [Ot] A. Otwinowska, *Monodromie d'une famille d'hypersurfaces; application à l'étude des variétés de Hodge*, preprint
- [SaM] M.Saito *Mixed Hodge modules*, Publ. RIMS, Kyoto Univ., **26** (1990), 221–333
- [Sch] P. Schneider, *Introduction to the Beilinson conjectures*, in: Beilinson's Conjectures on Special Values of L -functions (M. Rapoport, N. Schappacher and P. Schneider, ed), Perspectives in Math., **4**, Academic Press
- [T] T. Terasoma *Complete intersections with middle Picard number 1 defined over $\mathbb{Q}$* , Math. Z., **189** (1985), 289–296
- [V] C. Voisin, *Une précision concernant le théorème de Noether*, Math. Ann., **280** (1988), 605-611
- [SGA4] P. Deligne et al. *Théorie des Topos et Cohomologie Etale des Schémas*, Lecture Notes in Math., **305** (1970)